



# A note on a sum theorem for dimension $\mathcal{K}$ -Ind

Vitaly V. Fedorchuk<sup>1</sup>

Moscow State University (M.V. Lomonosov), Department of General Topology and Geometry, Moscow, Russia

## ARTICLE INFO

MSC:  
54F45

**Keywords:**  
Dimension  
Simplicial complex  
Hereditarily normal space  
Perfectly normal space  
Sum theorem

## ABSTRACT

Main results are:

1. Let  $Y$  be a closed subspace of a hereditarily normal  $X$  such that  $\mathcal{K}\text{-Ind } Y \leq n$  and  $\mathcal{K}\text{-Ind}(X \setminus Y) \leq n$ . Then  $\mathcal{K}\text{-Ind } X \leq n$ .
2. Let  $X$  be a perfectly normal space. Then a finite sum theorem for dimension  $\mathcal{K}\text{-Ind}$  holds in  $X$  if and only if  $\mathcal{K}\text{-Ind}$  is monotonic in  $X$ .

We denote by  $\mathcal{K}$  a non-empty set of finite complete simplicial complexes.

© 2011 Published by Elsevier B.V.

## 1. Introduction

In [4] inductive dimension function  $\mathcal{K}\text{-Ind}$ , where  $\mathcal{K}$  is a non-empty set of finite simplicial complexes, was introduced (look at Definition 2.4). This dimension is an extension of the classical inductive dimension  $\text{Ind}$ , since  $\{0, 1\}\text{-Ind } X = \text{Ind } X$  for every normal space. Generally,

$$\mathcal{K}\text{-Ind } X \leq \text{Ind } X.$$

In [4] it was proved that

$$\mathcal{K}\text{-Ind } X = \text{Ind } X$$

if and only if  $\mathcal{K}$  contains a disconnected complex  $K$ .

One of the main questions concerning dimension  $\mathcal{K}\text{-Ind}$  is the following one:

Let a perfectly normal space  $X$  be the union of its closed subspaces  $X_i$ ,  $i = 1, 2, \dots$ . Is it true that

$$\mathcal{K}\text{-Ind } X = \sup\{\mathcal{K}\text{-Ind } X_i; i = 1, 2, \dots\}?$$

The answer is unknown even if  $X = X_1 \cup X_2$ .

Here we prove (Theorem 3.4) that the finite sum theorem for dimension  $\mathcal{K}\text{-Ind}$  holds for subspaces of a perfectly normal space  $X$  if and only if dimension  $\mathcal{K}\text{-Ind}$  is monotonic in subspaces of  $X$ . The proof is based on the following

**Finite Dowker theorem** (Theorem 3.1). *Let  $Y$  be a closed subspace of a hereditarily normal space  $X$  such that  $\mathcal{K}\text{-Ind } Y \leq n$ ,  $\mathcal{K}\text{-Ind}(X \setminus Y) \leq n$ . Then  $\mathcal{K}\text{-Ind } X \leq n$ .*

E-mail address: [vvfedorchuk@gmail.com](mailto:vvfedorchuk@gmail.com).

<sup>1</sup> The author was supported by the Russian Foundation for the Basic research (Grant 09-01-00741) and the Program "Development of the Scientific Potential of Higher School" of the Ministry for Education of the Russian Federation (Grant 2.1.1.3704).

Let us recall that C.H. Dowker proved (look at [1,2]) the following

**Theorem D.** Let a hereditarily normal space  $X$  be the union of its subspaces  $X_i$ ,  $i = 1, 2, \dots$ , such that  $\text{Ind } X_i \leq n$ ,  $i = 1, 2, \dots$ , and  $\bigcup\{X_i: i = 1, 2, \dots, k\}$  is closed for  $k = 1, 2, \dots$ . Then  $\text{Ind } X \leq n$ .

Theorem 3.1 implies a finite version of Theorem D for dimension  $\mathcal{K}\text{-Ind}$ .

## 2. Preliminaries

**2.1.** In what follows  $\mathcal{K}$  stands for a non-empty set of finite complete simplicial complexes  $K$ , which we call *complexes*. For a complex  $K$  by  $v(K)$  we denote the set of all its vertices. A simplicial complex, which is the *nerve* of a finite family  $\alpha = \{A_1, \dots, A_s\}$  of sets, is denoted by  $N(\alpha)$ .

By a *space* we mean a topological normal  $T_1$ -space. For a space  $X$  by  $\exp X$  we denote the set of all closed subsets of  $X$ . By  $\text{Fin}_s(\exp X)$  we denote the set of all finite sequences  $\Phi = (F_1, \dots, F_m)$ ,  $F_j \in \exp X$ ,  $j = 1, \dots, m$ .

**Definition 2.2.** ([3]) Let  $X$  be a space,  $K$  be a complex, and  $\Phi = (F_1, \dots, F_m) \in \text{Fin}_s(\exp X)$ . A sequence  $u = (U_1, \dots, U_s)$ ,  $s \geq m$ , of open subsets of  $X$  is said to be a  $K$ -neighbourhood of  $\Phi$  if  $F_j \subset U_j$ ,  $j = 1, \dots, m$ , and there is an embedding  $N(u) \subset K$ . One can number vertices  $a_j \in v(K)$  so that the embedding  $N(u) \subset K$  is defined by the correspondence  $U_j \rightarrow a_j$ .

**Definition 2.3.** ([3]) A set  $P \subset X$  is called a  $K$ -partition of  $\Phi \in \text{Fin}_s(\exp X)$  (notation:  $P \in \text{Part}(\Phi, K)$ ) if  $P = X \setminus \bigcup u$ , where  $u$  is a  $K$ -neighbourhood of  $\Phi$ .

If a  $K$ -partition of  $\Phi$  exists, then  $N(\Phi) \subset K$ . Put

$$\text{Exp}_K(X) = \{\Phi \in \text{Fin}_s(\exp X): N(\Phi) \subset K\}. \quad (2.1)$$

**Definition 2.4.** ([4]) To every space  $X$  one assigns the dimension  $\mathcal{K}\text{-Ind } X$  which is an integer  $\geq -1$  or  $\infty$ . The dimension function  $\mathcal{K}\text{-Ind}$  is defined in the following way:

- (1)  $\mathcal{K}\text{-Ind } X = -1 \iff X = \emptyset$ ;
- (2)  $\mathcal{K}\text{-Ind } X \leq n$ , where  $n = 0, 1, \dots$ , if for every  $K \in \mathcal{K}$  and  $\Phi \in \text{Exp}_K(X)$  there exists a partition  $P \in \text{Part}(\Phi, K)$  such that  $\mathcal{K}\text{-Ind } P \leq n - 1$ ;
- (3)  $\mathcal{K}\text{-Ind } X = \infty$ , if  $\mathcal{K}\text{-Ind } X > n$  for all  $n \geq -1$ .

If the set  $\mathcal{K}$  contains only one complex  $K$ , we write  $\mathcal{K} = K$  and  $\mathcal{K}\text{-Ind } X = K\text{-Ind } X$ .

**Theorem 2.5.** For every space  $X$ ,  $\{0, 1\}\text{-Ind } X = \text{Ind } X$ .  $\square$

**Theorem 2.6.** ([4]) If  $Y$  is a closed subspace of a space  $X$ , then  $\mathcal{K}\text{-Ind } Y \leq \mathcal{K}\text{-Ind } X$ .  $\square$

**Theorem 2.7.** ([4]) If  $X = \bigoplus\{X_\alpha: \alpha \in A\}$  is a discrete union of spaces  $X_\alpha$ , then

$$\mathcal{K}\text{-Ind } X = \sup\{\mathcal{K}\text{-Ind } X_\alpha: \alpha \in A\}. \quad \square$$

**Lemma 2.8.** If  $U$  is an open  $F_\sigma$ -subset of a space  $X$ , then  $U$  is a cozero-set, i.e. there exists a continuous function  $\varphi: X \rightarrow [0, 1]$  such that  $U = \varphi^{-1}(0, 1]$ .  $\square$

**Strong swelling lemma 2.9.** Let  $\Phi = (F_1, \dots, F_m) \in \text{Fin}_s(\exp X)$ . Then there exists a family  $u = (U_1, \dots, U_m)$  of open subsets of  $X$  such that  $F_j \subset U_j$ ,  $j = 1, \dots, m$ , and  $N(\text{Cl}(u)) = N(\Phi)$ , where  $\text{Cl}(u) = (\text{Cl}(U_1), \dots, \text{Cl}(U_m))$ .  $\square$

**Nerve lemma 2.10.** ([4]) Let  $Y$  be subspace of a space  $X$ ,  $\alpha = (A_1, \dots, A_m)$  be a sequence of subsets of  $X$ , and  $\beta = (B_1, \dots, B_m)$  be a sequence of subsets of  $Y$  such that  $N(\alpha), N(\beta) \subset K$  and  $A_j \cap Y \subset B_j$ ,  $j = 1, \dots, m$ . Let  $C_j = A_j \cup B_j$  and  $\gamma = (C_1, \dots, C_m)$ . Then  $N(\gamma) \subset K$ .  $\square$

## 3. Main results

Let  $\mathcal{K}$  be a non-empty set of complexes and let  $X$  be a hereditarily normal space.

**Theorem 3.1.** Let  $Y$  be a closed subspace of a hereditarily normal space  $X$  such that  $\mathcal{K}\text{-Ind } Y \leq n$ ,  $\mathcal{K}\text{-Ind}(X \setminus Y) \leq n$ . Then  $\mathcal{K}\text{-Ind } X \leq n$ .

**Proof.** We shall apply induction with respect to  $n$ . For  $n = -1$  the theorem is obvious. Assume that the corresponding statements hold for dimensions less than  $n \geq 0$  and consider a hereditarily normal space  $X$  satisfying the assumption of our theorem. Let  $K \in \mathcal{K}$ ,  $\Phi = (F_1, \dots, F_m) \in \text{Exp}_K(X)$ , and  $\Phi|Y = (F_1 \cap Y, \dots, F_m \cap Y)$ . Then  $\Phi|Y \in \text{Exp}_K(Y)$ . Since  $\mathcal{K}\text{-Ind } Y \leq n$ , there is a family  $u = (U_1, \dots, U_m)$  of open subsets of  $Y$  such that

$$F_j \cap Y \subset U_j, \quad j = 1, \dots, m; \quad (3.1)$$

$$N(u) \subset K; \quad (3.2)$$

$$\mathcal{K}\text{-Ind}(Y \setminus U_1 \cup \dots \cup U_m) \leq n - 1. \quad (3.3)$$

Put  $P = Y \setminus U_1 \cup \dots \cup U_m$  and  $Z = X \setminus P$ . The family  $u$  is an open cover of a normal space  $Y \setminus P$ . Hence there exist closed subsets  $A_j$  of a space  $Y \setminus P$  such that

$$F_j \cap Y \subset A_j \subset U_j; \quad (3.4)$$

$$A_1 \cup \dots \cup A_m = Y \setminus P. \quad (3.5)$$

Put  $\alpha = (A_1, \dots, A_m)$ . From (3.2) and (3.4) it follows that

$$N(\alpha) \subset K. \quad (3.6)$$

Since  $Y \setminus P$  is closed in  $Z$ , the sets  $B_j = A_j \cup F_j$ ,  $j = 1, \dots, m$ , are closed in  $Z$ . Put  $\beta = (B_1, \dots, B_m)$ . The condition  $\Phi \in \text{Exp}_K(X)$  is equivalent to

$$N(\Phi) \subset K. \quad (3.7)$$

From (3.6), (3.7) and the Nerve lemma (Lemma 2.10) it follows that

$$N(\beta) \subset K. \quad (3.8)$$

Consequently, according to the Strong swelling lemma (Lemma 2.9) there exists a family  $v = (V_1, \dots, V_m)$  of open subsets of  $Z$  such that

$$B_j \subset V_j, \quad j = 1, \dots, m; \quad (3.9)$$

$$N(\delta) = N(\beta) \subset K, \quad (3.10)$$

where  $\delta = (D_1, \dots, D_m)$  and  $D_j = \text{Cl}_Z(V_j)$ .

Put  $E_j = D_j \setminus Y$ ,  $j = 1, \dots, m$ , and  $\epsilon = (E_1, \dots, E_m)$ . The sets  $E_j$  are closed in  $X \setminus Y$  and

$$N(\epsilon) \subset K \quad (3.11)$$

according to (3.10). But  $\mathcal{K}\text{-Ind}(X \setminus Y) \leq n$ . Consequently, according to (3.11) there exists a family  $w = (W_1, \dots, W_m)$  of open subsets of  $X \setminus Y$  such that

$$E_j \subset W_j, \quad j = 1, \dots, m; \quad (3.12)$$

$$N(w) \subset K; \quad (3.13)$$

$$\mathcal{K}\text{-Ind } Q \leq n - 1, \quad (3.14)$$

where

$$Q = X \setminus Y \cup W_1 \cup \dots \cup W_m. \quad (3.15)$$

Put  $G_j = V_j \cup W_j$ ,  $j = 1, \dots, m$ , and  $\gamma = (G_1, \dots, G_m)$ .

From (3.10), (3.13), and the Nerve lemma (Lemma 2.10) it follows that

$$N(\gamma) \subset K. \quad (3.16)$$

Condition (3.9) implies that

$$F_j \subset G_j, \quad j = 1, \dots, m. \quad (3.17)$$

Consequently,  $\gamma$  is a  $K$ -neighbourhood of  $\Phi$  in  $X$ . Then the set

$$R = X \setminus G_1 \cup \dots \cup G_m \quad (3.18)$$

is a  $K$ -partition of  $\Phi$  in  $X$ . We claim that

$$R = P \cup Q. \quad (3.19)$$

To check (3.19) it suffices to prove that

$$R \cap Y = P; \quad (3.20)$$

$$R \cap (X \setminus Y) = Q. \quad (3.21)$$

From definition of  $G_j$  it follows that

$$G_j \cap Y = V_j \cap Y. \quad (3.22)$$

Hence

$$(G_1 \cup \dots \cup G_m) \cap Y = (V_1 \cup \dots \cup V_m) \cap Y.$$

Condition (3.22) is equivalent to

$$Y \setminus G_1 \cup \dots \cup G_m = Y \setminus V_1 \cup \dots \cup V_m. \quad (3.23)$$

From (3.18) it follows that

$$Y \setminus G_1 \cup \dots \cup G_m = R \cap Y. \quad (3.24)$$

On the other hand,  $V_j \subset Z$  implies that  $(V_1 \cup \dots \cup V_m) \cap P = \emptyset$ . Consequently,  $P \subset Y \setminus V_1 \cup \dots \cup V_m \subset$  (in view of (3.9))  $Y \setminus B_1 \cup \dots \cup B_m \subset$  (because of  $B_j = A_j \cup F_j$ )  $Y \setminus A_1 \cup \dots \cup A_m =$  (in accordance with (3.5))  $= P$ .

Hence

$$P = Y \setminus V_1 \cup \dots \cup V_m. \quad (3.25)$$

From (3.24) and (3.25) we get (3.20).

The definition of  $E_j$  implies that

$$V_j \setminus Y \subset E_j, \quad j = 1, \dots, m. \quad (3.26)$$

Conditions (3.12) and (3.26) yield

$$V_j \setminus Y \subset W_j, \quad j = 1, \dots, m. \quad (3.27)$$

Consequently, the definition of  $G_j$  implies that

$$G_j \cap (X \setminus Y) = W_j. \quad (3.28)$$

Then  $R \cap (X \setminus Y) =$  (according to (3.18))  $= (X \setminus G_1 \cup \dots \cup G_m) \cap (X \setminus Y) = X \setminus Y \cup G_1 \cup \dots \cup G_m =$  (in view of (3.28))  $= X \setminus Y \cup W_1 \cup \dots \cup W_m =$  (because of (3.15))  $= Q$ .

Thus, the condition (3.21) is checked as well. Hence the equality (3.19) is proved. Since  $P \cap Q = \emptyset$ , we have  $Q = R \setminus P$ . On the other hand,

$$\mathcal{K}\text{-Ind } P \leq (\text{in view of (3.3)}) \leq n - 1;$$

$$\mathcal{K}\text{-Ind } Q \leq (\text{because of (3.14)}) \leq n - 1.$$

Consequently, by the inductive assumption we have

$$\mathcal{K}\text{-Ind } R \leq n - 1. \quad (3.29)$$

The condition (3.29) implies that  $\mathcal{K}\text{-Ind } X \leq n$ .  $\square$

Let us consider the following properties of a space  $X$ :

$(\mu_n)$  For each subspace  $Y \subset X$  and every open subspace  $U$  of  $Y$ , if  $\mathcal{K}\text{-Ind } Y \leq n$ , then  $\mathcal{K}\text{-Ind } U \leq n$ .

$(\mu_n^0)$  For each subspace  $Y \subset X$  and every open  $F_\sigma$ -subspace  $U$  of  $Y$ , if  $\mathcal{K}\text{-Ind } Y \leq n$ , then  $\mathcal{K}\text{-Ind } U \leq n$ .

$(\sigma_n)$  For each subspace  $Y \subset X$  and every pair  $Y_1, Y_2$  of closed subspaces of  $Y$  such that  $Y = Y_1 \cup Y_2$ , if  $\mathcal{K}\text{-Ind } Y_i \leq n$ ,  $i = 1, 2$ , then  $\mathcal{K}\text{-Ind } Y \leq n$ .

As a corollary of Theorem 3.1 we have

**Proposition 3.2.** *If a hereditarily normal space  $X$  has property  $(\mu_n)$ , then it also has property  $(\sigma_n)$ .*

**Proof.** Consider a subspace  $Y \subset X$  and a pair  $Y_1, Y_2$  of closed subspaces of  $Y$  such that  $Y = Y_1 \cup Y_2$  and  $\mathcal{K}\text{-Ind } Y_i \leq n$ ,  $i = 1, 2$ . By virtue of  $(\mu_n)$  the set  $Y \setminus Y_1$  satisfies the inequality  $\mathcal{K}\text{-Ind}(Y \setminus Y_1) \leq n$ . Applying Theorem 3.1 to the space  $Y$  and the pair  $Y_1, Y \setminus Y_1$  we obtain the inequality  $\mathcal{K}\text{-Ind } Y \leq n$ .  $\square$

**Proposition 3.3.** *If a hereditarily normal space  $X$  has property  $(\sigma_n)$ , then it also has property  $(\mu_n^0)$ .*

**Proof.** Let  $Y \subset X$  and let  $U$  be an open  $F_\sigma$ -set in  $Y$ . Then there exists a continuous function  $f : Y \rightarrow I$  such that  $U = f^{-1}((0, 1))$  (see Lemma 2.8). The sets  $B_i = f^{-1}([1/i + 1, 1/i])$ ,  $i = 1, 2, \dots$ , are closed in  $Y$ . By the closed subspace theorem (Theorem 2.6) we have

$$\mathcal{K}\text{-Ind } B_i \leq n, \quad i = 1, 2, \dots \quad (3.30)$$

Consider the sequences

$$B_{2i+1}, \quad i = 0, 1, 2, \dots; \quad B_{2i+2}, \quad i = 0, 1, 2, \dots$$

They are discrete. Put

$$A_1 = \bigcup \{B_{2i+1}, i = 0, 1, 2, \dots\}; \quad A_2 = \bigcup \{B_{2i+2}, i = 0, 1, 2, \dots\}.$$

By the Discrete sum theorem (Theorem 2.7) and (3.30) we have

$$\mathcal{K}\text{-Ind } A_1 \leq n; \quad \mathcal{K}\text{-Ind } A_2 \leq n.$$

But  $A_1 \cup A_2 = U$ . Consequently, property  $(\sigma_n)$  yields  $\mathcal{K}\text{-Ind } U \leq n$ .  $\square$

Since properties  $(\mu_n)$  and  $(\mu_n^0)$  are equivalent in perfectly normal spaces, from Propositions 3.2 and 3.3 we get

**Theorem 3.4.** *Properties  $(\mu_n)$  and  $(\sigma_n)$  are equivalent in the class of perfectly normal spaces.*  $\square$

**Question 3.5.** Does a perfectly normal space  $X$  satisfy property  $(\sigma_n)$ ,  $n = 0, 1, 2, \dots$ , for an arbitrary  $\mathcal{K}$ ?

**Remark 3.6.** The answer is “yes” if  $\mathcal{K}$  contains a disconnected complex  $K$ . In fact, in this case,  $\mathcal{K}\text{-Ind } X = \text{Ind } X$  (look at [4]) for every normal space  $X$ , and dimension  $\text{Ind}$  satisfies the countable sum theorem in the class of all perfectly normal spaces (look at [1,2]).

## References

- [1] C.H. Dowker, Inductive dimension of completely normal spaces, Quart. J. Math. Oxford 4 (1953) 267–281.
- [2] R. Engelking, Theory of Dimensions. Finite and Infinite, Sigma Ser. Pure Math., vol. 10, Heldermann, Lemgo, 1995.
- [3] V.V. Fedorchuk, Spaces which are weakly infinite-dimensional modulo simplicial complexes, Vestnik Moskov. Univ. Ser. I Mat. Mekh. 3 (2009) 33–40.
- [4] V.V. Fedorchuk, Inductive dimensions modulo simplicial complexes and ANR-compacta, Colloq. Math. 120 (2) (2010) 223–247.